



THE TASMANIAN NATURALIST

ISSN 0819-6826

POSTAL ADDRESS: G.P.O. BOX 68A, HOBART 7001

EDITOR: D.G. HIRD

A NEW RECORD FOR *BANKSIA SERRATA* IN TASMANIA

by *Phil Cullen*

Dept. Parks, Wildlife and Heritage, Tasmania

Banksia serrata (saw banksia) was probably first recorded in Tasmania by James Backhouse in the early 1830's at Sisters Beach on the North West Coast. Until recently this was the only known population in the state. However, information from Mr Alf Stackhouse of Killiecrankie, Flinders Island has led to the location of a stand of the species in the Wingaroo area on northern Flinders Island. He and his family had known of this occurrence for the past 40 years but had not publicised the fact. Here *B. serrata* forms an open woodland of about 1 ha on an undisturbed hillslope in the southern part of the new Wingaroo Nature Reserve. Given the known age of the stand (greater than 40 years), its location, and the size of the largest individuals (up to 8m tall), it is very unlikely that this population has developed from an introduction by European settlers.

B. serrata is widespread along the east coast of Australia from Fraser Island to Wilsons Promontory (Fig. 1) where it occurs in coastal and near coastal woodlands, forests and heathlands (Taylor and Hopper 1988). The species extends into the Blue Mountains on the mid-coast of New South Wales.

B. serrata belongs to an interesting group of plants which are widespread on the mainland but have very limited occurrences in this state, usually on islands in Bass Strait or at restricted localities along the north and east coasts of Tasmania. These species conform to three generalized distribution patterns. Some have strong affinities with temperate and warm temperate communities on the coastal plain of eastern and southeastern Australia. Examples include *B. serrata*, *B. integrifolia*, *Melaleuca armillaris*, *Leucopogon esquamatus*, *Helichrysium argophyllum*, *Sicyos australis*, and *Eleocharpos reticulatus*. A second subgroup is comprised of species which have affinities with semiarid mallee and arid desert

communities of western Victoria, South Australia, and Western Australia, for example *Eutaxia microphylla*, *Pimelea serpyllifolia*, *Geococcus pusillus*, *Lavatera plebia*, *Lasiopetalum discolor*, and *Myoporum parvifolium*. The remainder of the species are more widespread in southern and south eastern Australia, for example *Hakea sericia*, *H. ulicina*, and *Helichrysum baxteri*.

Why do these taxa have such limited distributions in Tasmania? Either, they are restricted by unfavourable climatic conditions or other factors, perhaps having been more widespread in the past, or else they are relatively recent arrivals at their present locations by long distance seed dispersal from the mainland but as yet have not colonised their potential ranges. The second explanation cannot be discounted for any of these taxa unless evidence is found which can establish their antiquity in our region. However, considerable congruence in the distribution patterns of the species suggests that they have experienced similar histories rather than being the survivors of unique dispersals across Bass Strait brought about by a fortuitous interplay of events.

It is possible that at least some of these species were widespread on the Bassian land bridge which connected Tasmania to the Mainland during the last glacial period when sea levels were up to 150 metres below present. Since then climate change and rising sea levels may have forced these taxa into refugial situations at sites where environmental factors remain marginally suitable for their continued existence.

The floras of Curtis and Rodondo island in Bass Strait contain elements which suggest that they are remnants from the Bassian Plain (Kirkpatrick et. al. 1974). One of these species, *Melaleuca armillaris*, is recorded from both these islands and also from Long Island near Cape Barren Island (Steve Harris pers. com.). *M. armillaris* is found in coastal communities throughout eastern Australia. There is a disjunction of over 300km between the Bass Strait populations and the nearest population in East Gippsland. The absence of *M. armillaris* from most islands in eastern Bass Strait suggests that the species is poorly dispersed across sea barriers. Disjunctions in the distribution of *B. serrata* (Fig. 1) are comparable to those of *M. armillaris*. Despite many apparently suitable habitats in northern Tasmania, and the islands, the species is virtually absent from the region. Coast banksia, *Banksia integrifolia*, has a similarly puzzling distribution. It occurs from south of Cooktown to near Geelong on the Australian coast and in two tiny populations in Bass Strait, King Island (now extinct) and Long Island in the Hogan Group (3 Individuals). Kirkpatrick et. al. (1974) raise the possibility that, during the last glacial, lower sea levels exposing a coastal plain may have allowed presently disjunct populations of *M. armillaris* to have been continuous with mainland populations but question whether climatic conditions at the time could allow such a distribution to occur. Pollen evidence from cave sediments on

Hunter Island (Hope 1978) indicates that the vegetation of the Bassian Plain, at the time of maximum glaciation, was a very open grassland community with composite shrubs or annuals and scattered eucalypts. The climate in the region would have been colder, drier, and windier than at present. Hope suggests that the Bassian vegetation would have had affinities with present day semi-arid, warm-temperate woodlands, and with inverted treeline and alpine timberline communities of the driest part of Tasmania's Central Plateau. Disjunct mallee and desert species found on the Bass Strait Islands provide supportive evidence for this view.

How then could species which are characterised by eastern temperate and warm temperate coastal climates survive the rigors of the Bassian Plain? Two answers to this question can be suggested. One, that the eastern margin of the Bassian Plain remained warm and moist enough to support coastal communities containing these species. This may have been facilitated by the moderating influence of warm east Australian ocean currents on the east coast of the Bassian Plain in the same manner that the Gulf Stream moderates the climate in north western Europe. Two, that following the intensely cold and dry glacial maximum increasing temperatures and precipitation in the late Pleistocene-early Holocene may have allowed these taxa to expand onto the Bassian Plain from refugia in far eastern Victoria before rising seas severed Tasmania and the island from the Mainland. The early Holocene climate for the region is considered to have been warmer and wetter than at present (Macphail 1980). Either way it appears that sea level rises and deterioration of climate in the middle Holocene have combined to dramatically limit the distribution of temperate/warm-temperate species like *B. serrata* in our region.

REFERENCES

- Hope, G.S. 1978 The late Pleistocene and Holocene vegetational history of Hunter Island, north-western Tasmania. *Aust. J. Bot.* 26:493-514.
- Kirkpatrick, J.B., Massey, J.S. and Parsons, R.S. 1974 Natural history of Curtis Island, Bass Strait. 2. Soils and vegetation with notes on Rodondo Island. *Pap. Proc. Roy. Soc. Tas.* 107:131-144
- Macphail, M.K. 1980 Regeneration processes in Tasmanian forests: A long-term perspective based on Pollen Analysis. *Search* 11:184-189.
- Taylor, A. and Hopper, S. 1988 The Banksia Atlas. Australian Flora and Fauna series No 8. Aust. Gov. Pub. Service.




Figure 1 The distribution of *Banksia serrata*
(updated from Taylor and Hooper (1988))